

E. Yu. Zyryanova¹, E. D. Kazakova¹, A. A. Musikhina^{1,2},
A. D. Atlasheva¹, O. N. Chupakhin^{1,2},
I. A. Utepova^{1,2}, V. A. Ishkhanyan¹

¹*Institute of Chemical Engineering, Ural Federal University,
620002, Russia, Ekaterinburg, Mira St., 28,
a.a.musikhina@urfu.ru,*


²*I. Ya. Postovsky Institute of Organic Synthesis,
Ural Branch of the Russian Academy of Sciences,
620108, Russia, Ekaterinburg, S. Kovalevskoy St., 22*

SYNTHESIS OF 1-AZINYL-1'-ISOPROPENYLFERROCENES*

Keywords: isopropenylferrocene, regioselective synthesis, Friedel–Crafts acylation, Wittig reaction.

Appended olefins in ferrocene system allow ones to create the high conjugated two-dimensional p-electron system [1] giving the wide scope of useful materials [2]. Moreover, metal-containing polymers appear to act in the processes of a redox-responsive release from patchy nanocapsules [3], catalysis modulating activity [4] and ion-selective membrane-gating through nanopores. The very fact of having a lipophilic ferrocene moiety in the structure of drugs modulates the range of their physiological activity while enhancing it [5].

In this work we combine olefins and azines fragments on the ferrocene matrix by regioselective acetylation of azines followed by the Wittig reaction.



Scheme 1. *Reagents and conditions:* i, Ac₂O, AlCl₃, CH₂Cl₂, 0 °C;
ii, Ph₃P=CH₂ (from Ph₃PMe⁺I⁻ and BuLi), Et₂O, rt.

* *This study was supported by the Russian Science Foundation (grant № 19-73-00243) and the Russian Foundation for Basic Research (Project № 20-43-660054).*